

Small Ground Mammal Habitat Associations in Wombat State Forest

P.V. Macak

2012

Arthur Rylah Institute for Environmental Research

Unpublished client report to the Wombat Biodiversity Working Group/Wombat Forestcare Inc.



Small ground mammal habitat associations in Wombat State Forest

Phoebe Macak

Arthur Rylah Institute for Environmental Research
123 Brown Street, Heidelberg, Victoria 3084

June 2012

Arthur Rylah Institute for Environmental Research
Department of Sustainability and Environment
Heidelberg, Victoria

Report produced by: Arthur Rylah Institute for Environmental Research
Department of Sustainability and Environment
PO Box 137
Heidelberg, Victoria 3084
Phone (03) 9450 8600
Website: www.dse.vic.gov.au/ari

© State of Victoria, Department of Sustainability and Environment 2012

This publication is copyright. Apart from fair dealing for the purposes of private study, research, criticism or review as permitted under the *Copyright Act 1968*, no part may be reproduced, copied, transmitted in any form or by any means (electronic, mechanical or graphic) without the prior written permission of the State of Victoria, Department of Sustainability and Environment. All requests and enquiries should be directed to the Customer Service Centre, 136 186 or email customer.service@dse.vic.gov.au

Citation: Macak, P.V. (2012) Small ground mammal habitat associations in Wombat State Forest. Arthur Rylah Institute for Environmental Research Unpublished client report to the Wombat Biodiversity Working Group/Wombat Forest Care Inc. Department of Sustainability and Environment, Heidelberg, Victoria

Disclaimer: This publication may be of assistance to you but the State of Victoria and its employees do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.

Accessibility:

If you would like to receive this publication in an accessible format, such as large print or audio, please telephone 136 186, or through the National Relay Service (NRS) using a modem or textphone/teletypewriter (TTY) by dialling 1800 555 677, or email customer.service@dse.vic.gov.au

Front cover photo: Left: community member (Viola) installing a ground-based hair-tube (Photo by Tibor Hegedis); Right: understorey wattle from a study site (Photo by Phoebe Macak)

Contents

List of tables and figures.....	vi
Acknowledgements.....	vii
Summary	1
1 Introduction.....	2
2 Methods.....	4
2.1 Study area	4
2.2 Site selection	4
2.3 Small ground mammal hair-tube surveys	4
2.4 Vegetation surveys.....	5
2.5 Data analyses	6
3 Results.....	7
3.1 Ground mammals.....	7
3.2 Vegetation surveys.....	8
3.3 Cover/abundance of vegetation attributes.....	9
3.4 Relationship between small ground mammals and habitat attributes	11
4 Discussion	13
References	15
Appendix 1 Study sites in Wombat State Forest used to investigate mammal habitat associations	16
Appendix 2 Tree form classification system used during eucalypt tree surveys	18
Appendix 3 Size and form of eucalypt trees measured at study sites	19
Appendix 4 Cover-abundance of understorey vegetation at each study site	21

List of tables and figures

List of tables

Table 1. Mammals recorded from hair collected during hair-tubes surveys within Wombat State Forest..... 8

Table 2. Number of trees counted per site, grouped into eucalypt species or type, size class and tree form. Dominate eucalypts are MM=Messmate or Gum=gum-barked 9

Table 3. Cover abundance measurements for habitat attributes and numbers of logs and stumps at each study site. 11

Table 4. Parameter estimates from best fit logistic regression model of the presence of Agile Antechinus in Wombat State Forest using data collected in 2005-2006..... 12

List of figures

Figure 1. Study sites within Wombat State Forest used to investigate mammal habitat associations in mature forest 3

Acknowledgements

The work presented in this report is an extension of a study on arboreal mammals and their habitat (results of which are presented in an earlier report) and was conducted at the same sites. Funding was provided by the Wombat Biodiversity Working Group (BWG) via the Wombat Community Forest Management Council of Stewards and the Department of Sustainability and Environment (DSE).

BWG and Wombat State Forest community members conducted the vegetation and hair-tube surveys (including hair-tube construction). Hair analysis was carried out by Barbara Triggs. Gayle Osborne and Tanya Loos (BWG) coordinated the data collection and collation.

Ryan Chick (ARI) provided the BWG with detailed instructions regarding construction and deployment of hair-tubes.

Statistical analysis was carried out by the author with advice from Michael Scroggie (ARI).

Richard Loyn and Matt Bruce provided comments on earlier drafts.

Note that the BWG has since been disbanded with the above members now part of Wombat Forestcare Inc.

Hair-tubing was conducted under Wildlife Act 1975 and National Parks Act 1975 Research Permit No. 10003522, and Department of Sustainability and Environment Arthur Rylah Institute for Environmental Research Animal Ethics Committee Approval No. 05/011.

Summary

Wombat State Forest is a large (70,000 ha) mixed eucalypt foothill forest that has had a long history of timber harvesting. Messmate *Eucalyptus obliqua* has been the major sawlog extracted. Sawlog harvesting ceased in 2003 and a forest management trial was initiated to continue forest management as a community-government partnership. The Wombat Forest Biodiversity Working Group (BWG) obtained funding to conduct research into aspects of forest biodiversity. The Arthur Rylah Institute for Environmental Research (ARI) was asked to carry out a study to contribute to the understanding of the fauna-habitat associations in mature forest areas of Wombat State Forest.

Thirty research sites of 1 ha each were selected in areas dominated by Messmate or gum-barked eucalypts. The BWG collected vegetation data (tree species and diameter classes, and cover abundance of understorey habitat attributes), and conducted hair-tubing for ground mammals using tree-mounted and ground-based tubes.

A total of eight mammal species (six native and two introduced) were recorded during spotlighting and hair-tubing. The species detected are known to occur in the area and are considered common in Victoria. Agile Antechinus *Antechinus agilis* was the most widespread species recorded (19 sites) and Swamp Rat *Rattus lutreolus* the least widespread (1 site). A total of 3,978 eucalypt trees over 40cm in diameter (DBH) were recorded with Messmate the most common, followed by peppermints, gum-barked eucalypts and stags. The number of trees at each site ranged from 78 to 178. The vegetation surveys revealed that several habitat attributes were present at most or all sites (including bracken, native grasses, rushes/sedges, shrub layer and *Acacia spp.*) while some were less common (including tree-ferns and Cherry Ballart *Exocarpus cupressiformis*): a summary of the results is provided.

Statistical analysis focused on the most common small mammals detected (Agile Antechinus *Antechinus agilis* and Bush Rat *Rattus fuscipes*), and tree and vegetation attributes. Logistic regression models found a positive relationship between increased cover of leaf litter and the presence of Agile Antechinus. No relationship was detected between the Bush Rat and habitat attributes.

Note that the survey effort used to collect small mammal data may not have been sufficient to detect all species at each site, or to identify patterns of occurrence in terms of habitat attributes.

ARI also conducted spotlighting surveys for arboreal mammals, the results of which were presented in an earlier report.

1 Introduction

Wombat State Forest (WSF), west-central Victoria, is characterised by mixed eucalypt foothill forest and covers an area of about 70,000 ha. It has a long history of timber harvesting, with Messmate *Eucalyptus obliqua* the major sawlog extracted. Widespread harvesting ceased in 2002 pending a review, and in 2003 a community forest management (CFM) trial was initiated, with the intention of forest management continuing as a community-government partnership. The last sawlog coupe was harvested in 2005. The last remaining sawlog licence was bought back by the State government in 2006. Since then there has been limited extraction of commercial and domestic firewood, wood chop logs, and timber for other products such as posts and poles.

The Wombat CFM Council of Stewards was formed to facilitate and coordinate community involvement in the ongoing management of the forest, overseeing a number of community working groups that addressed particular issues. In 2004 one of these groups, the Wombat Forest Biodiversity Working Group (BWG) obtained funding from the Department of Sustainability and Environment (DSE) through the Council of Stewards, to conduct research into a number of topics relating to biodiversity in WSF. The Arthur Rylah Institute for Environmental Research (ARI), DSE was asked to be involved and as a result a short study was proposed that would increase the understanding of fauna-habitat associations in mature forest areas of WSF.

The Wombat CFM Council of Stewards and associated working groups were disbanded in 2006. The community group Wombat Forestcare Incorporated was subsequently formed, effectively replacing the BWG, and has since become one of the major contributors to community liaison with DSE on forest management.

This report focuses on the relationships between small ground mammals and vegetation characteristics. It forms part of a larger study that investigated the relationships between mammals and vegetation characteristics at 30 sites across the forest (Figure 1). Part of the aim of the broad study was to assess the variation within stands classed as mature forest, and to determine the contributions made by different variables to habitat for fauna. A companion report related the presence of Greater Gliders (and to a lesser extent other arboreal mammals) and the number and proportion of broad types of eucalypt trees (i.e. stringybarks such as Messmates, peppermints and gum-barked species) (Macak *et al.* 2010). On request by the former BWG, the opportunity was taken to extend the study and increase knowledge about the distribution of small ground mammals over a wide geographic area of the forest, and to identify any obvious associations between the presence of these mammals and vegetation characteristics (habitat elements) of sites. Although many of the small mammals recorded from WSF are regarded as common, the survey method employed (hair-tubing) is also an effective way to identify cryptic and rare mammals that may not otherwise be easily detected.

Some of the information used here was initially presented in Macak *et al.* 2010 but has been included to enable this to be a stand-alone report.




Figure 1. Study sites within Wombat State Forest used to investigate mammal habitat associations in mature forest

2 Methods

2.1 Study area

Wombat State Forest is located about 70 km north-west of Melbourne, between Ballan and Woodend, and north of Daylesford to Yandoit, covering an area of about 70,000 ha. It is characterised by mixed species foothill forest, with Messmate the most common eucalypt. Other eucalypts present include Narrow-leaved Peppermint *Eucalyptus radiata*, Broad-leaved Peppermint *E. dives*, Manna Gum *E. viminalis*, Swamp Gum *E. ovata*, Mountain Grey Gum *E. cypellocarpa* and Candlebark *E. rubida*, with the last four species being gum-barked eucalypts. Study sites were selected from an area of the forest roughly encompassed by Daylesford, Spring Hill, North Blackwood, Blackwood, Blakeville and Barkstead.

2.2 Site selection

Thirty study sites were selected in large blocks of continuous mature forest dominated by Messmate or gum-barked species (Figure 1). Twenty-six of these sites had been used previously during other ARI research projects and were selected based on State Forest Resource and Inventory Project forest polygon codes that identify age class and dominant eucalypt species. The sites selected were all classified as mature forest. Including these sites in the current study reduced the amount of time spent on site selection and field reconnaissance to check access and suitability. Four additional sites were chosen from a selection made by the BWG. These were selected based on forest maturity, dominant eucalypt species, access and connectivity with the wider forest area. Sites were located a minimum of 1 km apart from each other to ensure that individual animals targeted during surveys would not be recorded from two sites on the same night. Sites were chosen to cover a large geographic area within the forest in order to sample a wide variety of habitats present across the landscape.

Each study site was set up as a 250 m x 40 m (1 ha) area. A 250 m transect line was measured along a compass bearing using a hipchain and marked every 10 m using flagging tape. The site extended 20 m on either side of the transect line. Coordinates (MGA GDA94) for each site were taken at the start of each transect (i.e. at 0 m) using a Garmin eTrex Vista global positioning system unit. Most sites (28) were located within the general management zone of the forest. Two sites were within the grounds of the Bullarto Reservoir.

2.3 Small ground mammal hair-tube surveys

The presence of small ground mammals was investigated by using hair-tubes (sampling devices designed to collect mammal hair). Two types of hair-tubes were used: small tree-mounted tubes and ground-based tunnels. The two types were arranged alternately along the transect line of each site every 10 m equalling a total of 24 tubes per site (12 of each type). Tubes were left *in situ* for 14–16 days between October–December 2005. The hair-tube survey was conducted in two stages, with half (15) the sites surveyed at a time. Tubes were collected, processed for hair, cleaned and re-taped between the two stages. Sites were selected within each survey stage to cover the geographic spread of the study area, and sites closest to each other were not surveyed at the same time. The topographic position of each tube was estimated (ridge, upper slope, mid slope, lower slope, drainage line, gully, flat). A grand total of 720 hair-tubes (360 each of tree and ground) were deployed over the 30 study sites. Mammal hair caught on the sticky surface of the adhesive tape was identified to species level, where possible. Hair samples were analysed by Barbara Triggs ('Dead Finish', Genoa, Victoria)

Hair-tubes were constructed and deployed by the BWG after instructions from Ryan Chick (ARI). All hair-tubes were baited with a mixture of honey, peanut butter and rolled oats.

Tree-mounted tubes were constructed from PVC piping (38 mm diameter x 120 mm long). At each end of the piping three strips of double-sided adhesive tape were stuck over the edge such that a strip extended for about 40 mm lengthwise along the outside and inside surface of the piping. Bait was contained within a small PVC tube (about 36 mm long) and held in place in the middle of the hair-tube when attached to the trunks of the largest trees (favouring eucalypts) by a nail that went through the diameter of the piping and down the length of the bait holder.

Ground-based tunnels were made from clear polycarbonate sheets (250 mm x 300 mm) ('Handiglaze Premium'; Laserlite Australia, Cheltenham Vic.). A stainless steel double-spoon type tea infuser was used to contain bait, and was placed into a 22 mm circular hole cut 100 mm along the long edge and 12.5 cm along the short edge of the polycarbonate sheet. Four pieces of double sided adhesive tape (48 mm wide x 120 mm long) were placed along the length of the short edge of the sheet (two on each edge) over four pieces of reinforced garden hose (600 mm long). The sheets were bent and held in place in the ground by a length of galvanised wire (5 mm diameter x 600 mm long) bent into a 'U' shape. The wire passed through the protruding handle of the baitholder to hold the components together. The tunnels were placed with the short edge along the ground.

One site (WCR28) was re-surveyed using tree-mounted tubes during February 2007, in an effort to detect Brush-tailed Phascogales *Phascogale tapoatafa*. Three parallel lines of 12 tubes each (a total of 36 tubes) were deployed. Tubes were set at 20 m apart; the lines were 200 m from each other. Honey-water was sprayed onto the bark above and below tree-mounted tubes to enhance their attractiveness. These results will be presented separately to those from the main study.

2.4 Vegetation surveys

Vegetation surveys were designed to give an indication of the presence of structural and habitat elements of the forest, particularly those that provide shelter or foraging substrate for arboreal and small ground-based mammals. Some of these elements were chosen with the view that any future work on diurnal birds could draw on the survey results. Vegetation surveys were carried out by members of the BWG during 2006.

All live and dead (stag) eucalypts greater than or equal to 40cm diameter at breast height (DBH) were measured with a diameter tape and classified into three size ranges (40-79, 80-99, >100 cm DBH). Trees were also categorised into one of eight tree forms according to a scale representing increasing senescence (Appendix 2). Live trees were identified to species level in the case of Messmates or grouped according to their bark type as either peppermints or gum-barked eucalypts. The bark grouping was used as some peppermints and gum-barked eucalypts are difficult to identify to species level. Species included within the peppermint grouping included Narrow-leaved Peppermint and Broad-leaved Peppermint. Gum-barked eucalypts included Manna Gum, Mountain Grey Gum, Swamp Gum and Candlebark.

To collect data on understorey trees, shrubs, logs and stumps the site was divided into 13 segments along the length of the transect. Each segment was 40 m wide, with 12 segments 20 m long (e.g. 0-20 m, 20-40 m etc.) and one segment 10 m long (the end of each transect i.e. 240-250 m). The density of understorey trees and shrubs was rated by visual assessment according to the Braun-Blanquet scale of cover abundance; * = rare, 1 = <5%, 2 = 5-25%, 3 = 25-50%, 4 = 50-75%, 5 = 75-100%. Cover abundance surveys were carried out by the same two observers at all sites. Understorey trees and shrubs were recorded by species or in broad groupings as follows: Blackwood *Acacia melanoxylon*, Silver Wattle *A. dealbata*, Cherry Ballart *Exocarpus cupressiformis*, Musk Daisy-bush *Olearia argophylla*, Coprosma *Coprosma spp.*, woody shrubs < 2 m tall, woody shrubs >2 m tall, sedges, ground ferns, tree-ferns and grasses. Note that the woody shrub grouping did not include Blackwood, Silver Wattle, Musk Daisy-bush or Coprosma.

The number of logs > 40 cm DBH within each site segment were categorised as 40-59 cm, 60-79 cm or >80 cm DBH. The DBH of logs was measured at the widest part of the log. Where a log crossed into two segments, it was only included in the segment embracing the longest part of the log. Hence for some segments, the number of logs present may be an underestimate. The number of stumps >40 cm DBH within each site segment was recorded and placed into one of three size classes: 40-59 cm, 60-79 cm or >80 cm DBH. The DBH for logs was measured at the top of stumps. For the purposes of summarising vegetation data to allow comparisons between sites, information for each segment was averaged post-survey to give an overall cover abundance rating for each site.

2.5 Data analyses

Data were analysed to explore the relationships between the presence of ground mammals recorded during hair-tube surveys and combinations of tree variables and the cover/abundance of vegetation attributes (numbers of individual trees, both live trees and stags; numbers of trees and tree types within DBH classes; numbers of logs and stumps within DBH classes; cover of shrubs, ferns and leaf litter). Overall cover abundances calculated post-survey were used. Analyses were undertaken using simple logistic regressions (generalised linear models). Akaike's information criterion (corrected for small sample size) (AICc) was calculated for each model to allow them to be ranked for best fit, with the lowest AICc being the most suitable one. AICc differences and Akaike weights were then used to further assess which model(s) best fit the data (Burham and Anderson 2002). Only the best-fit model is presented.

3 Results

3.1 Ground mammals

A total of eight mammal species were recorded from hair-tubes during the study (Table 1): Agile Antechinus *Antechinus agilis*, Bush Rat *Rattus fuscipes*, Swamp Rat *Rattus lutreolus*, Brushtail Possum *Trichosurus* sp., Common Wombat *Vombatus ursinus*, Swamp Wallaby *Wallabia bicolor*, House Cat *Felis catus* and Red Fox *Vulpes vulpes*. For a more comprehensive list of all fauna recorded during the study (including during spotlighting surveys) see Macak *et al.* 2010. The most common species recorded were Agile Antechinus (19 sites), Swamp Wallaby (15 sites) and Brushtail Possums (11 sites). The Swamp Rat was recorded from one site only. Two species of Brushtail Possum have been recorded in Wombat State Forest (Common Brushtail Possum *T. vulpecula* and Mountain Brushtail Possum *T. cunninghami*) and they cannot be distinguished reliably on the basis of hair samples. Hence all records were considered as “Brushtail Possums”. Although many sites recorded unidentified antechinus (including at site 4 where this was the only antechinus record), only confirmed records of Agile Antechinus are included here and were used for analysis.

All species except Agile Antechinus were recorded from ground installed tubes only. Agile Antechinus were recorded mostly from tree-mounted tubes (90% of records), and at only one site was it recorded solely from a ground-tube and not from tree-mounted tubes.

The additional hair-tubing session at site WCR28 was carried out when hair analysis from the main round of surveys detected ‘probable’ Brush-tailed Phascogale from two tree-mounted tubes. All 36 tubes deployed during the second hair-tube survey returned hair identified as Agile Antechinus. Due to the ‘probable’ status of the Brush-tailed Phascogale identification, the presence of this species at this site remains unconfirmed.

Table 1. Mammals recorded from hair collected during hair-tubes surveys within Wombat State Forest.

Site	Agile Antechinus	Bush Rat	Swamp Rat	Unidentified Brushtail Possum	Common Wombat	Swamp Wallaby	House Cat	Red Fox
WCR01	X					X		X
WCR02	X			X		X		
WCR03	X							
WCR04	^	X			X	X		
WCR05	X	X		X				
WCR06							X	
WCR07	X					X		
WCR08	X							
WCR09						X		
WCR10								
WCR11		X		X				
WCR12	X	X				X		
WCR13	X			X			X	
WCR14	X					X		X
WCR15	X					X		
WCR16						X		
WCR17	X							
WCR18	X							
WCR19	X			X	X			
WCR20	X			X				
WCR21	X	X				X		
WCR22	X							
WCR23	X	X		X				
WCR24						X		
WCR25	X	X						
WCR26				X		X		
WCR27				X		X		
WCR28	X			X	X	X		
WCR29			X	X	X	X		
WCR30		X						
Total	19	8	1	11	4	15	2	2

^ unidentified antechinus

3.2 Vegetation surveys

A total of 3,978 eucalypt trees were identified and classified into three size classes during the surveys (Table 2). Messmate was the most common eucalypt (2,301) followed by peppermints (920), gum-barked eucalypts (611) and stags (146). The number of trees at each site ranged from 78 to 178. Peppermints were the most widespread, present at 29 sites. Messmates were present at all but three sites, while there were ten sites with no gum-barked eucalypts recorded.

Most trees recorded were within the smallest DBH class 40-79 cm (3,310), followed by 80-99 cm (545) and >100 cm (123). Trees in the 40-79 cm and 80-99 cm DBH range were present at all sites, while three sites had no trees >100 cm DBH. Only three sites had more than ten trees >100 cm DBH, with 19 being the maximum (WCR05).

The majority of stags were classified as forms 7 and 8, the most decayed in the scale of tree senescence when trees are effectively a solid or hollow stump. Stags were recorded from all but one site.

Table 2. Number of trees counted per site, grouped into eucalypt species or type, size class and tree form. Dominate eucalypts are MM=Messmate or Gum=gum-barked

Site	Dominate eucalypt*	Number of trees	Gum-barked	Messmate	Peppermint	Stag	40-79cm DBH	80-99cm DBH	>100cm DBH	Total live trees	Form 1	Form 2	Form 3	Form 4	Form 5	Form 6	Form 7	Form 8
WCR01	MM	163	7	152	3	1	116	41	6	162	162	0	1	0	0	0	0	0
WCR02	MM	178	1	159	18	0	159	19	0	178	175	3	0	0	0	0	0	0
WCR03	MM	147	0	117	27	3	138	8	1	144	143	1	1	0	0	0	2	0
WCR04	MM	142	0	114	27	1	124	17	1	141	137	4	0	0	0	0	0	1
WCR05	MM	132	3	91	35	3	84	29	19	129	125	4	0	1	0	0	0	2
WCR06	MM	136	36	79	16	5	119	13	4	131	115	16	1	0	0	0	0	4
WCR07	MM	124	12	108	2	2	112	12	0	122	121	1	0	0	0	0	0	2
WCR08	MM	90	24	65	0	1	67	22	1	89	89	0	0	0	0	0	0	1
WCR09	MM	130	6	105	5	14	103	23	4	116	108	8	5	3	1	1	1	3
WCR10	MM	122	0	106	10	6	108	12	2	116	116	0	1	0	0	0	0	5
WCR11	MM	128	0	95	25	8	111	15	2	120	107	13	1	0	0	0	0	7
WCR12	MM	149	0	98	41	10	144	5	0	139	139	0	0	0	1	3	3	3
WCR13	MM	142	2	66	71	3	121	17	4	139	138	1	0	0	0	1	1	1
WCR14	MM	102	0	53	45	4	77	20	5	98	94	4	0	0	0	0	4	0
WCR15	MM	127	9	33	74	11	112	13	2	116	90	26	0	1	1	0	6	3
WCR16	MM	174	0	123	44	7	160	12	2	167	145	22	0	0	0	0	1	6
WCR17	MM	150	0	131	16	3	120	17	13	147	138	9	0	0	0	0	2	1
WCR18	MM	157	1	134	19	3	116	38	3	154	143	11	0	0	0	0	3	0
WCR19	Gum	112	28	39	42	3	90	15	7	109	105	4	0	0	0	0	1	2
WCR20	Gum	99	20	0	74	5	81	13	5	94	84	10	2	0	0	0	2	1
WCR21	Gum	91	60	1	24	6	75	12	4	85	82	3	1	0	0	3	2	0
WCR22	Gum	149	43	0	99	7	130	15	4	142	136	6	0	0	0	0	3	4
WCR23	Gum	121	66	0	51	4	107	12	2	117	111	6	1	0	0	0	2	1
WCR24	Gum	78	60	5	9	4	51	21	6	74	67	7	0	4	0	0	0	0
WCR25	MM	168	0	140	19	9	154	11	3	159	159	0	2	1	0	0	1	5
WCR26	Gum	104	59	14	26	5	68	25	11	99	91	8	0	0	0	0	1	4
WCR27	Gum	148	102	1	37	8	122	24	2	140	135	5	0	2	1	5	0	0
WCR28	MM	129	4	107	16	2	89	34	6	127	124	3	1	0	0	1	0	0
WCR29	Gum	148	68	45	30	5	135	11	2	143	136	7	0	0	0	1	0	4
WCR30	MM	138	0	120	15	3	117	19	2	135	133	2	1	1	0	1	0	0
Total		3978	611	2301	920	146	3310	545	123	3832	3648	184	18	13	4	16	35	60
Mean		132.6	20.4	76.7	30.7	4.9	110.3	18.2	4.1	127.7	121.6	6.1	0.6	0.4	0.1	0.5	1.2	2.0
Std Dev		25.3	28.1	51.7	24.0	3.3	27.8	8.5	4.1	25.4	26.1	6.4	1.0	1.0	0.3	1.2	1.5	2.1

3.3 Cover/abundance of vegetation attributes

Overall, the study sites showed a small degree of variation in terms of shrubs and tree species present (Table 3). Mistletoe was present at seven sites, and only rarely at those. Of the two *Acacia* species surveyed, most sites had one or the other, sometimes both. Blackwood was recorded from more sites than Silver Wattle. Blackwood was present mainly in shrub form with 1-

3 canopy-sized individuals recorded from 11 sites. When present, *Acacia* was usually rare throughout sites, in low densities (up to 5% cover). Cherry Ballart was present at three sites in low densities. Musk Daisy-bush was recorded at eight sites from scattered individuals up to 5% cover, with two large trees (in the 40-60 cm DBH range) at WCR09. All sites had a shrub layer, except for WCR06, where a few scattered Acacias were the only shrubs. Coprosma was present at many sites, at low to very low densities. All sites had varying cover of leaf litter, generally between 5-50%, with three sites recording over 50%. Bracken and native grasses were present at all sites, generally in low densities, although a few sites had a high cover of grasses. Rushes/sedges were present at all sites, usually in low densities. Tree-ferns were scarce across the sites, present at eight, while ground ferns were recorded from the majority of sites, but at low densities. The number of logs and stumps varied between sites. Logs within the 40-59 cm DBH range were present at all sites and were more numerous than those within the larger DBH ranges 60-79 cm and >80 cm, which were not always present. The total number of logs at individual sites ranged from 1 to 31, with 355 logs recorded overall. Stumps were present at all sites, with those in the smaller size range (40-59 cm) more numerous than larger sized stumps. The total number of stumps counted per site ranged from 4 to 35, with 503 stumps recorded overall.

Within sites, the cover abundance of some vegetation attributes varied across the 13 segments while some were spread evenly throughout the 1ha area (Appendix 4). For example, rushes/sedges were sometimes concentrated in high densities over several adjacent segments, while being sparse within the rest of the site. This change in density over the transect sometimes corresponded to the landform at different points of a site with the highest densities of rushes/sedges recorded in gullies and drainage lines.

Table 3. Cover abundance measurements for habitat attributes and numbers of logs and stumps at each study site. Cover abundance values are averages of 13 measurements taken at each site according to the Braun-Blanquet scale: *=rare; 1=<5%; 2=5-25%; 3=25-49%; 4=50-75%; 5=75-100%.

Site	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree-ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Total logs	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm	Total stumps
WCR01	0	0	*	0	1	1	1	1	3	2	2	2	0	0	3	1	1	5	10	8	3	21
WCR02	0	0	0	0	0	2	2	0	3	2	2	1	0	0	5	1	1	7	18	7	2	27
WCR03	0	0	*	0	0	1	2	*	3	1	3	1	0	*	10	3	3	16	7	6	3	16
WCR04	0	0	*	0	0	1	1	0	1	2	4	1	0	0	3	0	0	3	8	4	2	14
WCR05	0	0	*	0	0	1	1	0	2	2	3	2	0	1	16	0	0	16	31	2	2	35
WCR06	0	*	*	0	0	0	0	0	3	2	2	2	0	*	1	0	0	1	8	5	1	14
WCR07	0	0	*	0	0	0	1	0	4	2	1	1	0	*	6	0	0	6	9	5	0	14
WCR08	0	0	1	*	0	1	1	0	3	2	2	1	0	0	5	1	0	6	12	4	2	18
WCR09	*	0	*	0	1	2	1	1	1	2	4	1	*	*	6	2	2	10	14	3	1	18
WCR10	0	0	0	0	0	1	1	*	2	2	2	1	0	0	9	4	3	16	13	4	4	21
WCR11	0	0	0	0	0	2	2	*	2	2	3	1	*	*	10	3	1	14	9	3	3	15
WCR12	0	0	0	0	0	2	2	0	2	2	2	1	*	*	8	3	2	13	11	5	2	18
WCR13	0	0	*	0	*	2	1	0	2	1	2	1	0	*	8	2	1	11	16	8	1	25
WCR14	*	0	0	0	1	1	1	*	2	1	3	1	0	0	9	2	0	11	3	3	0	6
WCR15	1	1	0	0	0	1	2	0	3	1	1	1	0	0	13	2	0	15	10	0	0	10
WCR16	*	0	0	*	0	1	1	*	3	1	1	1	*	*	11	7	1	19	20	6	0	26
WCR17	0	0	0	*	0	1	1	0	4	2	1	1	0	1	21	7	3	31	6	5	4	15
WCR18	*	0	0	*	0	1	1	*	2	2	3	2	*	*	14	7	1	22	14	10	3	27
WCR19	*	1	*	0	0	1	1	0	2	2	2	2	0	0	21	1	2	24	6	7	5	18
WCR20	0	1	1	0	0	1	1	1	2	2	1	1	0	0	9	1	4	14	5	5	0	10
WCR21	0	1	*	0	0	1	*	0	3	2	2	1	0	*	4	3	5	12	12	2	0	14
WCR22	0	1	0	*	0	*	*	0	5	1	1	1	0	0	4	3	0	7	8	0	0	8
WCR23	0	1	1	0	0	1	1	0	3	1	2	1	0	*	6	0	1	7	4	1	0	5
WCR24	0	0	1	0	0	1	1	1	2	2	2	2	0	1	8	0	0	8	5	4	0	9
WCR25	0	0	0	0	1	1	1	1	2	1	2	2	*	1	7	5	2	14	8	9	2	19
WCR26	0	1	*	0	0	1	1	1	3	2	1	1	0	*	9	4	0	13	10	6	2	18
WCR27	0	*	*	0	0	1	2	*	2	1	2	3	0	1	5	1	0	6	10	0	0	10
WCR28	0	0	*	*	1	1	1	1	3	2	1	1	1	*	14	4	0	18	14	5	1	20
WCR29	0	0	*	0	*	2	2	1	2	2	2	3	0	*	5	0	0	5	3	1	0	4
WCR30	*	0	1	0	1	1	1	1	3	2	2	2	*	2	3	1	1	5	19	4	5	28

3.4 Relationship between small ground mammals and habitat attributes

Agile Antechinus were most commonly recorded at localities within sites where there were no Musk Daisy Bush, Silver Wattle, Blackwood or tree ferns, or where there was a low level of shrub density (up to 25% cover-abundance), bracken, rushes/sedges or ground ferns (Appendix 4).

There were usually 0-2 logs and 2-3 stumps close by. The Bush Rat, the next most common small mammal found, was recorded in hair-tube localities where there was bracken and rushes/sedges present (usually as a low cover/abundance), no Silver Wattle, no more than a sparse cover of Blackwood, a medium cover of shrubs, no or very few tree ferns and no or a low cover of ground

ferns. There were usually logs in the immediate vicinity (up to three) but very few stumps, if any. The only site that the Swamp Rat was detected was one of two that had the highest density of rushes/sedges (25-49%).

Statistical analysis was only carried out on the more common of the small mammals, Agile Antechinus and Bush Rat. Logistic regressions indicated that was a positive relationship between the presence of Agile Antechinus and an increasing cover of leaf litter (Table 4). No other relationships were detected between this species and other habitat attributes (cover of shrubs and ferns, numbers of logs and stumps, and numbers of trees) during statistical analysis. There was no detectable link between records of Agile Antechinus and the placement of tubes in the landscape.

No such relationships were detected for the Bush Rat and there were insufficient data for the Swamp Rat to undertake statistical analysis.

Table 4. Parameter estimates from best fit logistic regression model of the presence of Agile Antechinus in Wombat State Forest using data collected in 2005-2006.

Parameter	Estimate	Standard error	Test statistic	Probability
Constant	-2.09	1.47	-1.42	0.156
Cover of leaf litter	1.068	0.593	1.80	0.071
Model				0.043

4 Discussion

The three small mammal species detected during this study are known to be widespread in Victoria, including in Wombat State Forest (Menkhorst 1995, Atlas of Victorian Wildlife, DSE 2007). Many previous studies of forest mammals have shown that Bush Rats and Agile Antechinus are common in a range of habitats whereas Swamps Rats are more localised (Menkhorst 1995, Seebeck 1995a, Seebeck 1995b). This pattern of occurrence is echoed by the current study and reflects the habitat requirements of these species.

Agile Antechinus climb readily, nesting in tree hollows from ground level to the tops of trees and sometimes in fallen logs (Cockburn and Lazenby-Cohen 1992). They nest communally, requiring large hollows to do so, although females may nest solitarily at times (Cockburn and Lazenby-Cohen 1992). They are insectivorous, foraging in leaf litter, fissures in bark and tree trunks and, some studies suggest, from the indented surfaces of ferns, tree ferns and certain shrubs such as *Olearia* (Gullan and Robinson 1980). This species is very widespread and has been found in a wide variety of environments. It appears to be associated with a dense shrub layer (Dickman 1980, Lindenmayer *et al.* 1994, Cunningham *et al.* 2005); probably due to the cover it provides (Laidlaw and Wilson 2006) and/or the increased availability of foraging habitat (Gullan and Robinson 1980). However, not all studies have found such associations (Claridge *et al.* 2008). A high density of animals has been recorded at some sites with a large area of leaf litter cover (due to the litter providing a greater foraging substrate), while high densities have been also been recorded at sites with low cover of leaf litter (surmised as being a surrogate for a greater cover of shrubs) (Cunningham *et al.* 2005). In the current study leaf litter was shown to have a positive influence on the presence of Agile Antechinus, i.e. the higher the cover of litter, the more likely this species would be present at a site. Other factors that influence the presence of this species include the availability of suitable nesting hollows which are usually in stags or large mature trees (Cockburn and Lazenby-Cohen 1992). The presence of hollows was not assessed during the current study as there are many difficulties and potential errors associated with identifying hollows (Koch 2008). However, tree size data were collected and this attribute may give an indication of the incidence of hollows (Gibbons *et al.* 2000). No relationships between tree attributes and the presence of the Agile Antechinus were detected in the current study.

Bush Rats were far less commonly recorded across the study sites than Agile Antechinus, probably partly due to their slightly different habitat requirements. Bush Rats are known to prefer wet habitats and a structurally complex and dense shrubby understorey (Lindenmayer *et al.* 1994, Laidlaw and Wilson 2006) and none of the shrubs measured were present in high levels of cover/abundance. Rushes and sedges, indicators of a moist environment, were occasionally recorded in higher densities than shrubs, however this was only at two sites and Bush Rats were not detected at these sites.

In the current study there were a limited number of variables that could be tested together in predictive models as the relatively low number of sites (30) meant that models needed to be kept simple, with only a few variables included at a time. Many habitat elements were only recorded at a small number of sites, giving many absences in the data and making it difficult to build predictive models for those variables. For some variables, such as shrubs, there was little variation in the data and that inhibits the ability of the type of predictive models used here to detect a relationship between cover/abundance and fauna species presence.

Further studies on the small mammal component of Wombat State Forest may benefit from measuring variables that are targeted to known habitat preferences for particular species of interest. A more precise scale for cover/abundance, e.g. actual percentage cover, would give the

data more subtlety and a better chance of teasing out influences on the presence of those particular species.

References

- Claridge, A.W, Tennant, P., Chick, R. and Barry S.C. (2008) Factors influencing the occurrence of small ground-dwelling mammals in southeastern mainland Australia. *Journal of Mammalogy* **89(4)**: 916-923
- Cockburn, A. and Lazenby-Cohen, K.A. (1992) Use of nest trees by *Antechinus stuartii*, a semelparous lekking marsupial. *Journal of Zoology, London* **226(4)**: 657-680
- Cunningham, R.B., Lindenmayer, D.B., MacGregor, C., Barry, S. and Welsh, A. (2005) Effects of trap position, trap history, microhabitat and season on capture probabilities of small mammals in a wet eucalypt forest. *Wildlife Research* **32(8)**: 657–671
- Dickman, C.R. (1980) Ecological studies of *Antechinus stuartii* and *Antechinus flavipes* (Marsulialia: Dasyuridae) in open-forest and woodland habitats. *Australian Zoologist* **20(3)**: 433-446
- Gibbons, P., Lindenmayer, D.B., Barry, S.C. and Tanton, M.T. (2000) Hollow formation in eucalypts from temperate forests in southeastern Australia. *Pacific Conservation Biology* **6**: 218-228
- Gullan, P.K. and Robinson, A.C. (1980) Vegetation and small mammals of a Victorian forest. *Australian Mammalogy* **3**: 87-95
- Koch, A.J. (2008) Errors associated with two methods of assessing tree hollow occurrence and abundance in *Eucalyptus obliqua* forest, Tasmania. *Forest Ecology and Management* **255(3-4)**: 674-685
- Laidlaw, W.S., and Wilson, B.A. (2006) Habitat utilisation by small mammals in a coastal heathland exhibiting symptoms of *Phytophthora cinnamomi* infestation. *Wildlife Research* **33**: 639–649
- Lindenmayer, D.B., Cunningham, R.B., Donnelly, C.F., Triggs, B.E. and Belvedere, M. (1994) Factors influencing the occurrence of mammals in retained linear strips (wildlife corridors) and contiguous stands of montane ash forest in the Central Highlands of Victoria, southeastern Australia. *Forest Ecology and Management* **67**: 113-133
- Macak, P.V., Chick, R.R and Loyn, R.H. (2010) Arboreal mammal habitat associations in Wombat State Forest. Arthur Rylah Institute for Environmental Research Unpublished client report to the Wombat Biodiversity Working Group/Wombat Forestcare Inc. Department of Sustainability and Environment, Heidelberg
- Menkhorst, P.V. (1995) Brown Antichinus. Pp 44-47 *In* Menkhorst, P.W. (ed) Mammals of Victoria: distribution, ecology and conservation. Oxford University Press, Melbourne
- Seebeck, J.H. (1995) Bush Rat. Pp 224-226 *In* Menkhorst, P.W. (ed) Mammals of Victoria: distribution, ecology and conservation. Oxford University Press, Melbourne
- Seebeck, J.H. (1995) Swamp Rat. Pp 227-228 *In* Menkhorst, P.W. (ed) Mammals of Victoria: distribution, ecology and conservation. Oxford University Press, Melbourne

Appendix 1 Study sites in Wombat State Forest used to investigate mammal habitat associations

Site	Dominant eucalypt	Map sheet ¹ (1:25,000)	MGA easting [^]	MGA northing [^]	Compass bearing [°]	Location notes and directions*
WCR01	Messmate	Bullarto	251431	5850166	220	Cairns Road, 900m south of Farm Road, almost opposite Rat Hole Track.
WCR02	Messmate	Bullarto	251331	5851588	250	Farm Track, 450m north of Farm Road (take left fork 300m from gate). Track is 400m north-west of Cairns Road, and 2km east of Specimen Hill Road.
WCR03	Messmate	Bullarto	250205	5855215	0	Leonards Hill-Bullarto South Road, 700m west of Wombat Creek Rd - opposite water point.
WCR04	Messmate	Bullarto	253667	5858269	230	Site is ~50m up track off Osborne Road, track ~450m north of Trentham-Daylesford Road.
WCR05	Messmate	Bullarto	255852	5858430	230	site 100m up unmarked track off Babbington Rd, track 500m north of Daylesford-Trentham Road. Within Bullarto Reservoir area.
WCR06	Messmate	Cleever Hill	249313	5844590	80	Blakeville-Bunding Road, 200m east of Ballan-Daylesford Road.
WCR07	Messmate	Barkstead	243380	5847762	260	Barkstead South Road, 2.4km north of Spargo Creek Road & ~ 180m north of Broken Track, opposite dam.
WCR08	Messmate	Barkstead	243938	5846489	190	Barkstead South Road, 950m north of Spargo Creek Road.
WCR09	Messmate	Cleever Hill	255489	5844213	175	off track along Binks Rd pointing to a water point; site 100m along track, site crosses track. Track 300m east of Neville Track & 1.75km east of Blackville Road.
WCR10	Messmate	Greendale	260646	5845163	90	Charcoal Tk ~500m south of Greenhills Road. Track 600m north-east of Paradise Road.
WCR11	Messmate	Trentham	260658	5846763	50	Blackwood Ridge Road, ~750m west-north-west of Greenhills Road.
WCR12	Messmate	Bullarto	255868	5848921	20	Nolan Creek Road, 200m east of Stockyard Track.
WCR13	Messmate	Bullarto	255561	5850697	230	Nolan Creek Road, 2km north of Stockyard Track & 1.1km south of Lerderderg Road.
WCR14	Messmate	Trentham	261109	5852567	355	Yankee Road, 680m south-west of Pronks Track. NOTE old mine shafts present.
WCR15	Messmate	Trentham	267520	5849557	260	Ambler Lane ~ 950m south of Chettle Road (through Parks Vic road closure gate).
WCR16	Messmate	Trentham	267648	5850639	200	Chettle Road ~200m north-west of Ambler Lane. Along gully.


Site	Dominant eucalypt	Map sheet ¹ (1:25,000)	MGA easting [^]	MGA northing [^]	Compass bearing [°]	Location notes and directions*
WCR17	Messmate	Trentham	266169	5851207	250	Chettle Road, 2.9km south-east of Golden Point Road, 1.75km south-east of Wombat Track, (600m south of waterpoint).
WCR18	Messmate	Trentham	263407	5851542	250	Old Blackwood Road, 800m south of Yankee Road.
WCR19	gum-barked	Bullarto	256053	5859713	230	200m west along track off Babbington Hill Road. Track ~30m north of Top Track & 500m north of Bullarto Reservoir Road. Within Bullarto Reservoir area.
WCR20	gum-barked	Bullarto	249949	5846663	200	Track off McGee Road. Track ~360m north-west of Hell Hole Track. Site ~280m west of McGee Road.
WCR21	gum-barked	Coliban	261578	5869008	220	Track off Spring Hill Road (almost opposite another track) ~580m east of Mains Track. Site 200m south of SpringHill. Site crosses track. NOTE old mine shafts present.
WCR22	gum-barked	Coliban	260818	5867212	30	Chinaman Track 1.75km north of Mudlark Road. Transect crosses track. NOTE old mine shafts present.
WCR23	gum-barked	Coliban	257808	5861105	30	Springs Road, 100m south of Loddon River Road & 550m north of Mineral Springs Picnic Ground.
WCR24	gum-barked	Bullarto	249329	5851402	190	Unmarked track off Farm Road - just west of Specimen Hill Road intersection.
WCR25	Messmate	Bullarto	253984	5848184	160	Blackwood Ridge Road <100m east of Gentle Annie Track & Stockyard Track, 600m east of Camp Road
WCR26	gum-barked	Daylesford	255033	5862884	330	Rack north off Kangaroo Creek Road, 1.2km east of Osborne Road and 200m east of Spout Track.
WCR27	gum-barked	Trentham	259044	5858780	0	Blighs Lane/Road, 800m east of Coliban Drive
WCR28	Messmate	Trentham	264563	5851808	55	Golden Point Road, 1.7km south of Chettle Road.
WCR29	Messmate/ gum-barked	Bullarto	255345	5852274	120	Roach Road, 50m north of Lynch Track, just north of creek (east side of road).
WCR30	Messmate	Bullarto	251351	5856091	260	Wombat Creek Track, 1.25km from Leonards Hills-Bullarto South Road.

¹ Mapsheet numbers; Daylesford 7723-3-1, Coliban 7723-2-4, Barkstead 7723-3-3, Bullarto 7723-3-2, Trentham 7723-2-3, Cleever Hill 7722-4-1

[^] coordinates in GDA94, Zone 55

*more detailed directions are provided in Macak *et al.* 2010

Appendix 2 Tree form classification system used during eucalypt tree surveys



Classification of forms of trees (modified from Lindenmayer *et al.* 1991b). Tree form characteristics: 1. Living tree; 2. living tree with dead or broken top; 3. stag (dead tree) with most of branches still intact; 4. stag with 0-25% of the top broken off (branches remaining as stubs only); 5. stag with the top 25-50% broken away; 6. stag with the top 50-75% broken away; 7. a solid stag with > 75% of the top broken away; and 8. hollow stump.

Appendix 3 Size and form of eucalypt trees measured at study sites

Only trees over 40cm DBH were measured. Trees are grouped according to species or bark type

Eucalypts	Messmate										Peppermint										Gum-barked										Stag																		Grand Total
DBH class	40-59 cm			60-79 cm			>80 cm			Total	40-59 cm			60-79 cm			>80 cm			Total	40-59 cm			60-79 cm			>80 cm			Total	40-59 cm								60-79 cm						>80 cm			Total	
Site/Form	1	2	Total	1	2	Total	1	2	Total		1	2	Total	1	2	Total	1	2	Total		1	2	Total	1	2	Total	1	2	Total		3	4	5	6	7	8	Total	3	5	6	7	8	Total	7	8	Total			
WCR01	109		109	39		39	4		4	152	3		3							3	4		4	1		1	2		2	7								1					1				1	163	
WCR02	139	2	141	18		18				159	17		17		1	1				18	1		1							1																		178	
WCR03	107	1	108	8		8	1		1	117	27		27							27											1				2		3										3	147	
WCR04	95	3	98	16		16				114	25	1	26	1		1				27																								1	1	1	1	142	
WCR05	54		54	21		21	14	2	16	91	24	1	25	8		8	1	1	2	35	3		3							3		1				1	2								1	1	3	132	
WCR06	60	5	65	9	2	11	2	1	3	79	13	2	15					1	1	16	29	5	34	2		2				36	1					4	5									5	136		
WCR07	97		97	11		11				108	2		2							2	10	1	11	1		1				12						2	2									2	124		
WCR08	46		46	18		18	1		1	65											21		21	3		3				24										1	1				1	90			
WCR09	78	3	81	19	1	20	4		4	105	3	2	5							5	3	2	5	1		1				6	4	3		1	1	3	12	1	1				2			14	130		
WCR10	98		98	6		6	2		2	106	6		6	4		4				10											1					3	4				2	2				6	122		
WCR11	79	1	80	12	1	13	2		2	95	13	11	24	1		1				25																7	7	1				1			8	128			
WCR12	94		94	4		4				98	41		41							41														1	2	3	3	9			1			1			10	149	
WCR13	49		49	14		14	3		3	66	69	1	70	1		1				71	1		1	1		1				2					1		1			1		1	1	3	142				
WCR14	31		31	18		18	4		4	53	39	4	43	2		2				45															3		3					1		1	4	102			
WCR15	22	6	28	2	2	4		1	1	33	54	16	70	4		4				74	5		5	2	1	3	1		1	9		1	1		6	1	9					2	2				11	127	
WCR16	113	1	114	7	1	8	1		1	123	24	17	41		3	3				44															1	4	5					1	1		1	1	7	174	
WCR17	100	4	104	14	1	15	9	3	12	131	13	1	14	2		2				16															1	1	2					1		1	3	150			
WCR18	95	1	96	29	6	35	2	1	3	134	15	2	17	1	1	2				19				1		1				1					3		3								3	157			
WCR19	30		30	6	1	7	2		2	39	36	1	37	5		5				42	20	2	22	1		1	5		5	28					1		1					2	2				3	112	
WCR20											54	7	61	9	2	11	1	1	2	74	17		17	1		1	2		2	20	2				1		3				1		1		1	1	5	99	
WCR21	1		1							1	23		23	1		1				24	44	2	46	10		10	3	1	4	60	1			2	2		5			1						6	91		
WCR22											86	6	92	7		7				99	32		32	7		7	4		4	43					3	3	6					1	1				7	149	
WCR23											42	6	48	3		3				51	56		56	8		8	2		2	66	1				2		3					1	1				4	121	
WCR24	4		4	1		1				5	7	1	8				1		1	9	32	3	35	19	1	20	3	2	5	60		4					4									4	78		
WCR25	127		127	11		11	2		2	140	19		19							19												2	1			1	4	8							1	1	9	168	
WCR26	7		7	4		4	3		3	14	13	2	15	4	4	8	3		3	26	42		42	12	1	13	3	1	4	59					1	3	4							1	1	5	104		
WCR27				1		1				1	34	1	35	2		2				37	77	2	79	19	2	21	2		2	102		2	1	5			8									8	148		
WCR28	68	1	69	32		32	5	1	6	107	15		15		1	1				16	4		4							4	1					1			1				1			2	129		
WCR29	39	1	40	5		5				45	26	3	29	1		1				30	59	2	61	4	1	5	2		2	68				1		4	5								5	148			
WCR30	98	1	99	19		19	2		2	120	14	1	15							15												1	1		1			3								3	138		
Grand Total	1840	30	1870	344	15	359	63	9	72	2301	757	86	843	56	12	68	6	3	9	920	460	19	479	93	6	99	29	4	33	611	15	13	3	12	32	43	118	3	1	4	1	10	19	2	7	9	146	3978	
Mean	61.3	1.0	62.3	11.5	0.5	12.0	2.1	0.3	2.4	76.7	25.2	2.9	28.1	1.9	0.4	2.3	0.2	0.1	0.3	30.7	15.3	0.6	16.0	3.1	0.2	3.3	1.0	0.1	1.1	20.4	0.5	0.4	0.1	0.4	1.1	1.4	3.9	0.1	0.03	0.1	0.03	0.3	0.6	0.1	0.2	0.3	5.0	132.6	
Std Dev	44.0	1.6	44.3	10.1	1.2	10.4	3.0	0.7	3.6	51.7	20.4	4.5	22.4	2.6	1.0	2.9	0.6	0.3	0.7	24.0	21.5	1.2	22.1	5.4	0.5	5.7	1.4	0.4	1.7	28.1	0.9	1.0	0.3	1.0	1.4	1.9	3.1	0.3	0.2	0.3	0.2	0.7	0.7	0.3	0.4	0.5	3.2	25.3	

Appendix 4 Cover-abundance of understorey vegetation at each study site

Each site has data for 13 segments, plus an overall site average that was calculated post-survey. Cover-abundance was collected according to the Braun-Blanquet scale: * = rare, 1 = <5%, 2 = 5-25%, 3 = 25-50%, 4 = 50-75%, 5 = 75-100%

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR01	0-20m	0	0	0	0	0	*	*	*	4	2	1	1	0	0	0	0	0	1	0	0
WCR01	20-40m	0	0	0	0	0	*	*	*	4	2	1	*	0	0	0	0	0	0	2	0
WCR01	40-60m	0	0	0	0	0	1	*	*	3	2	1	1	0	0	0	0	1	1	0	1
WCR01	60-80m	0	0	0	0	*	1	*	*	2	2	1	1	0	0	0	0	0	0	1	0
WCR01	80-100m	0	0	*	0	*	*	1	1	2	2	1	2	0	0	1	0	0	0	0	0
WCR01	100-120m	0	0	0	0	0	1	1	*	2	2	2	2	0	0	1	0	0	0	0	0
WCR01	120-140m	0	0	*	0	0	1	*	1	2	2	2	2	0	0	0	0	0	2	0	1
WCR01	140-160m	0	0	0	0	1	1	1	1	2	2	2	2	0	0	0	0	0	0	1	0
WCR01	160-180m	0	0	0	0	*	1	1	1	2	2	2	2	0	0	0	0	0	1	1	0
WCR01	180-200m	0	0	0	0	*	*	1	1	3	2	1	2	0	0	0	0	0	2	1	0
WCR01	200-220m	0	0	0	0	1	1	1	1	3	1	2	1	0	0	0	0	0	0	0	0
WCR01	220-240m	0	0	0	0	1	1	1	1	3	2	2	1	0	0	0	0	0	3	0	1
WCR01	240-250m	0	0	0	0	*	1	1	1	3	2	2	1	0	0	1	1	0	0	2	0
WCR01	average	0	0	*	0	1	1	1	1	3	2	2	2	0	0	3	1	1	10	8	3
WCR02	0-20m	0	0	0	0	0	1	1	0	3	2	2	1	0	0	0	0	0	1	0	0
WCR02	20-40m	0	0	0	0	0	1	2	0	3	2	3	1	0	0	0	0	0	1	0	0
WCR02	40-60m	0	0	0	0	0	1	1	0	3	2	2	1	0	0	0	0	0	0	2	1
WCR02	60-80m	0	0	0	0	0	1	1	0	3	2	2	1	0	0	0	0	0	0	0	0
WCR02	80-100m	0	0	0	0	0	2	1	0	3	2	2	2	0	0	0	1	0	2	2	0
WCR02	100-120m	0	0	0	0	0	2	2	0	2	2	2	2	0	0	0	0	0	2	1	0
WCR02	120-140m	0	0	0	0	0	2	2	0	2	2	2	1	0	0	0	0	0	2	0	0
WCR02	140-160m	0	0	0	0	0	2	2	0	2	2	3	1	0	0	0	0	0	1	1	0
WCR02	160-180m	0	0	0	0	0	2	2	0	3	3	2	1	0	0	1	0	0	3	0	0
WCR02	180-200m	0	0	0	0	0	2	2	0	3	2	2	1	0	0	0	0	1	2	0	1
WCR02	200-220m	0	0	0	0	0	2	2	0	3	2	2	1	0	0	0	0	0	2	0	0
WCR02	220-240m	0	0	0	0	0	1	1	0	3	2	2	1	0	0	4	0	0	2	1	0
WCR02	240-250m	0	0	0	0	0	1	1	0	4	2	1	1	0	0	0	0	0	0	0	0
WCR02	average	0	0	0	0	0	2	2	0	3	2	2	1	0	0	5	1	1	18	7	2
WCR03	0-20m	0	0	0	0	0	1	3	0	2	1	3	1	0	*	0	0	0	2	0	0
WCR03	20-40m	0	0	0	0	0	1	3	*	1	1	3	1	0	0	0	0	0	0	0	0
WCR03	40-60m	0	0	0	0	0	1	3	0	2	1	3	1	0	0	0	0	0	0	0	0
WCR03	60-80m	0	0	0	0	0	*	1	0	2	1	3	1	0	0	1	0	0	2	0	0
WCR03	80-100m	0	0	*	0	0	*	*	0	3	2	3	1	0	0	0	1	2	0	0	2

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR03	100-120m	0	0	0	0	0	*	1	0	2	2	3	1	0	0	0	0	0	0	1	0
WCR03	120-140m	0	0	0	0	0	0	*	0	3	1	3	1	0	0	1	0	0	2	0	0
WCR03	140-160m	0	0	0	0	0	0	*	0	3	1	3	1	0	0	0	1	1	0	2	1
WCR03	160-180m	0	0	0	0	0	0	*	0	3	1	3	1	0	0	3	0	0	0	1	0
WCR03	180-200m	0	0	0	0	0	0	*	0	2	1	4	1	0	0	3	0	0	1	0	0
WCR03	200-220m	0	0	0	0	0	*	1	0	3	2	3	1	0	0	2	0	0	0	0	0
WCR03	220-240m	0	0	0	0	0	*	1	0	3	2	3	1	0	0	0	1	0	0	2	0
WCR03	240-250m	0	0	0	0	0	0	1	0	3	1	2	1	0	0	0	0	0	0	0	0
WCR03	average	0	0	*	0	0	1	2	*	3	1	3	1	0	*	10	3	3	7	6	3
WCR04	0-20m	0	0	*	0	0	*	*	0	1	2	4	1	0	0	0	0	0	1	1	0
WCR04	20-40m	0	0	*	0	0	1	1	0	1	1	4	1	0	0	0	0	0	0	0	0
WCR04	40-60m	0	0	*	0	0	1	1	0	1	2	5	1	0	0	0	0	0	0	0	0
WCR04	60-80m	0	0	0	0	0	1	1	0	1	1	4	1	0	0	0	0	0	0	0	0
WCR04	80-100m	0	0	0	0	0	1	1	0	1	1	5	1	0	0	0	0	0	0	0	0
WCR04	100-120m	0	0	*	0	0	1	1	0	1	2	5	1	0	0	1	0	0	1	0	1
WCR04	120-140m	0	0	1	0	0	1	1	0	1	2	5	1	0	0	0	0	0	1	0	0
WCR04	140-160m	0	0	0	0	0	1	1	0	1	1	5	2	0	0	1	0	0	1	1	0
WCR04	160-180m	0	0	0	0	0	1	1	0	1	1	5	1	0	0	0	0	0	1	0	0
WCR04	180-200m	0	0	0	0	0	*	1	0	1	1	4	1	0	0	0	0	0	1	0	1
WCR04	200-220m	0	0	0	0	0	1	1	0	1	2	4	1	0	0	1	0	0	1	0	0
WCR04	220-240m	0	0	*	0	0	1	1	0	3	1	3	1	0	0	0	0	0	0	2	0
WCR04	240-250m	0	0	0	0	0	1	1	0	4	1	1	1	0	0	0	0	0	1	0	0
WCR04	average	0	0	*	0	0	1	1	0	1	2	4	1	0	0	3	0	0	8	4	2
WCR05	0-20m	0	0	0	0	0	1	1	0	2	1	2	1	0	0	0	0	0	0	0	2
WCR05	20-40m	0	0	0	0	0	2	1	0	2	1	2	1	0	0	1	0	0	1	0	0
WCR05	40-60m	0	0	0	0	0	2	1	0	2	1	3	1	0	0	1	0	0	2	0	0
WCR05	60-80m	0	0	*	0	0	1	2	0	2	1	3	1	0	0	0	0	0	3	0	0
WCR05	80-100m	0	0	*	0	0	*	2	0	2	1	3	2	0	0	0	0	0	3	0	0
WCR05	100-120m	0	0	0	0	0	1	1	0	1	2	3	2	0	0	1	0	0	2	0	0
WCR05	120-140m	0	0	*	0	0	2	1	0	1	1	3	3	0	*	3	0	0	4	0	0
WCR05	140-160m	0	0	1	0	0	*	0	0	1	2	3	2	0	2	1	0	0	3	0	0
WCR05	160-180m	0	0	*	0	0	1	1	0	2	2	3	2	0	*	2	0	0	3	0	0
WCR05	180-200m	0	0	0	0	0	1	1	0	2	2	3	2	0	0	2	0	0	3	0	0
WCR05	200-220m	0	0	0	0	0	*	1	0	2	2	3	2	0	0	1	0	0	3	0	0
WCR05	220-240m	0	0	0	0	0	1	1	0	2	2	2	1	0	0	4	0	0	3	1	0
WCR05	240-250m	0	0	0	0	0	0	1	0	2	1	2	1	0	0	0	0	0	1	1	0
WCR05	average	0	0	*	0	0	1	1	0	2	2	3	2	0	1	16	0	0	31	2	2
WCR06	0-20m	0	*	0	0	0	0	0	0	5	1	1	1	0	*	0	0	0	0	0	0

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR06	20-40m	0	*	*	0	0	0	0	0	2	1	2	3	0	*	0	0	0	0	0	0
WCR06	40-60m	0	0	0	0	0	0	0	0	1	2	3	3	0	*	0	0	0	0	0	0
WCR06	60-80m	0	0	0	0	0	0	0	0	2	1	2	2	0	*	0	0	0	0	0	0
WCR06	80-100m	0	0	*	0	0	0	0	0	2	1	2	1	0	0	0	0	0	1	2	0
WCR06	100-120m	0	0	0	0	0	0	0	0	2	3	2	*	0	0	0	0	0	1	1	0
WCR06	120-140m	0	0	0	0	0	0	0	0	2	2	2	1	0	0	0	0	0	0	0	1
WCR06	140-160m	0	0	0	0	0	0	0	0	4	2	*	0	0	0	0	0	0	1	0	0
WCR06	160-180m	0	0	0	0	0	0	0	0	4	2	*	1	0	0	0	0	0	0	0	0
WCR06	180-200m	0	0	0	0	0	0	0	0	4	2	1	0	0	0	0	0	0	2	0	0
WCR06	200-220m	0	0	0	0	0	0	0	0	3	2	3	0	0	0	0	0	0	1	2	0
WCR06	220-240m	0	0	0	0	0	0	0	0	3	1	2	0	0	0	0	0	0	1	0	0
WCR06	240-250m	0	0	*	0	0	0	0	0	3	2	3	1	0	0	1	0	0	1	0	0
WCR06	average	0	*	*	0	0	0	0	0	3	2	2	2	0	*	1	0	0	8	5	1
WCR07	0-20m	0	0	*	0	0	0	*	0	4	2	1	1	0	0	1	0	0	1	0	0
WCR07	20-40m	0	0	0	0	0	0	*	0	4	2	1	1	0	0	0	0	0	1	0	0
WCR07	40-60m	0	0	0	0	0	0	*	0	4	1	1	1	0	0	0	0	0	0	0	0
WCR07	60-80m	0	0	0	0	0	0	*	0	4	1	1	1	0	0	1	0	0	1	0	0
WCR07	80-100m	0	0	0	0	0	0	*	0	4	1	1	1	0	0	1	0	0	1	0	0
WCR07	100-120m	0	0	0	0	0	0	*	0	4	1	1	1	0	0	0	0	0	0	0	0
WCR07	120-140m	0	0	0	0	0	0	*	0	4	1	1	1	0	1	1	0	0	1	0	0
WCR07	140-160m	0	0	0	0	0	0	1	0	4	2	1	1	0	1	0	0	0	0	3	0
WCR07	160-180m	0	0	0	0	0	0	1	0	4	2	1	1	0	0	0	0	0	0	2	0
WCR07	180-200m	0	0	0	0	0	0	1	0	4	2	1	1	0	0	0	0	0	0	0	0
WCR07	200-220m	0	0	1	0	0	0	1	0	4	2	1	1	0	0	0	0	0	1	0	0
WCR07	220-240m	0	0	1	0	0	0	1	0	4	2	1	2	0	0	2	0	0	2	0	0
WCR07	240-250m	0	0	1	0	0	0	1	0	4	2	1	2	0	0	0	0	0	1	0	0
WCR07	average	0	0	*	0	0	0	1	0	4	2	1	1	0	*	6	0	0	9	5	0
WCR08	0-20m	0	0	0	*	0	*	*	0	4	*	1	0	0	0	0	0	0	1	0	0
WCR08	20-40m	0	0	0	1	0	0	*	0	4	1	1	1	0	0	2	0	0	2	0	0
WCR08	40-60m	0	0	0	*	0	*	1	0	3	2	2	1	0	0	0	1	0	0	0	1
WCR08	60-80m	0	0	0	0	0	*	1	0	3	2	2	1	0	0	1	0	0	1	0	1
WCR08	80-100m	0	0	0	0	0	*	1	0	3	2	2	1	0	0	0	0	0	1	0	0
WCR08	100-120m	0	0	0	0	0	0	1	0	3	2	2	2	0	0	0	0	0	0	0	0
WCR08	120-140m	0	0	0	0	0	0	1	0	3	2	2	1	0	0	0	0	0	0	0	0
WCR08	140-160m	0	0	0	0	0	0	1	0	2	2	2	1	0	0	0	0	0	1	0	0
WCR08	160-180m	0	0	0	0	0	0	1	0	2	2	3	1	0	0	0	0	0	1	0	0
WCR08	180-200m	0	0	1	0	0	*	*	0	2	2	3	1	0	0	1	0	0	0	2	0
WCR08	200-220m	0	0	1	0	0	1	1	0	1	2	3	1	0	0	1	0	0	3	1	0

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR08	220-240m	0	0	1	0	0	1	1	0	1	3	3	1	0	0	0	0	0	2	0	0
WCR08	240-250m	0	0	1	0	0	0	1	0	2	3	2	2	0	0	0	0	0	0	1	0
WCR08	average	0	0	1	*	0	1	1	0	3	2	2	1	0	0	5	1	0	12	4	2
WCR09	0-20m	0	0	0	0	1	1	*	0	1	1	3	1	0	0	0	0	0	1	1	0
WCR09	20-40m	0	0	0	0	1	1	*	0	1	1	3	1	0	0	0	0	1	0	0	0
WCR09	40-60m	0	0	0	0	*	*	*	0	2	1	3	1	0	0	0	1	0	1	0	1
WCR09	60-80m	0	0	0	0	3	3	*	0	1	2	5	*	0	0	0	0	0	2	0	0
WCR09	80-100m	0	0	0	0	2	2	0	*	*	4	4	1	*	1	1	0	0	0	0	0
WCR09	100-120m	0	0	0	0	1	1	*	0	2	3	4	1	*	*	1	0	0	0	0	0
WCR09	120-140m	0	0	0	0	1	2	1	0	1	2	5	1	0	*	1	1	0	0	0	0
WCR09	140-160m	*	0	0	0	1	2	1	0	1	2	5	1	0	0	0	0	0	3	0	0
WCR09	160-180m	0	0	0	0	*	1	1	1	1	2	4	2	0	0	0	0	1	2	0	0
WCR09	180-200m	0	0	0	0	1	2	1	*	1	1	4	1	0	0	1	0	0	1	0	0
WCR09	200-220m	0	0	*	0	0	1	1	*	1	2	4	1	0	0	1	0	0	2	0	0
WCR09	220-240m	0	0	0	0	*	1	*	*	1	2	4	1	0	0	1	0	0	1	1	0
WCR09	240-250m	*	0	0	0	1	1	1	0	1	1	4	1	0	0	0	0	0	1	1	0
WCR09	average	*	0	*	0	1	2	1	1	1	2	4	1	*	*	6	2	2	14	3	1
WCR10	0-20m	0	0	0	0	0	1	3	0	2	1	2	1	0	0	1	1	0	2	0	0
WCR10	20-40m	0	0	0	0	0	1	2	0	3	1	1	1	0	0	0	1	0	0	0	0
WCR10	40-60m	0	0	0	0	0	1	1	0	4	2	1	1	0	0	0	0	0	4	1	0
WCR10	60-80m	0	0	0	0	0	1	1	0	3	2	2	1	0	0	1	0	0	0	0	0
WCR10	80-100m	0	0	0	0	0	0	0	0	3	2	3	1	0	0	2	0	0	0	0	0
WCR10	100-120m	0	0	0	0	0	*	*	0	3	2	2	1	0	0	1	0	2	0	0	1
WCR10	120-140m	0	0	0	0	0	*	*	0	2	3	2	1	0	0	1	0	0	1	0	0
WCR10	140-160m	0	0	0	0	0	2	*	0	2	2	2	1	0	0	0	0	0	1	1	0
WCR10	160-180m	0	0	0	0	0	*	*	*	2	2	2	1	0	0	1	0	0	0	1	0
WCR10	180-200m	0	0	0	0	0	1	*	*	2	2	2	1	0	0	0	1	1	0	1	1
WCR10	200-220m	0	0	0	0	0	1	*	0	2	1	2	1	0	0	1	1	0	0	0	1
WCR10	220-240m	0	0	0	0	0	1	*	*	2	2	2	1	0	0	1	0	0	4	0	1
WCR10	240-250m	0	0	0	0	0	1	*	*	2	1	2	1	0	0	0	0	0	1	0	0
WCR10	average	0	0	0	0	0	1	1	*	2	2	2	1	0	0	9	4	3	13	4	4
WCR11	0-20m	0	0	0	0	0	3	2	0	2	2	1	1	0	0	1	0	0	0	1	1
WCR11	20-40m	0	0	0	0	0	2	2	0	2	2	3	1	0	0	2	1	0	1	1	0
WCR11	40-60m	0	0	0	0	0	2	2	0	2	2	3	1	0	0	1	0	0	0	0	0
WCR11	60-80m	0	0	0	0	0	2	2	0	2	1	2	1	0	0	1	0	0	0	0	1
WCR11	80-100m	0	0	0	0	0	2	2	0	2	1	2	1	0	0	0	0	0	3	0	0
WCR11	100-120m	0	0	0	0	0	3	2	*	2	1	3	1	0	0	0	2	1	1	0	1
WCR11	120-140m	0	0	0	0	0	3	2	0	2	2	3	1	0	0	1	0	0	2	0	0

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR11	140-160m	0	0	0	0	0	3	3	0	2	2	3	2	*	*	2	0	0	1	0	0
WCR11	160-180m	0	0	0	0	0	3	3	0	2	2	3	1	0	0	0	0	0	0	0	0
WCR11	180-200m	0	0	0	0	0	2	2	0	2	2	3	1	*	0	0	0	0	0	0	0
WCR11	200-220m	0	0	0	0	0	2	2	0	2	2	3	1	0	0	1	0	0	1	1	0
WCR11	220-240m	0	0	0	0	0	2	2	0	2	1	3	1	*	0	1	0	0	0	0	0
WCR11	240-250m	0	0	0	0	0	1	2	0	2	1	3	1	0	0	0	0	0	0	0	0
WCR11	average	0	0	0	0	0	2	2	*	2	2	3	1	*	*	10	3	1	9	3	3
WCR12	0-20m	0	0	0	0	0	3	2	0	2	2	3	1	0	0	1	0	0	0	0	0
WCR12	20-40m	0	0	0	0	0	3	2	0	2	2	3	1	0	0	0	0	1	1	0	0
WCR12	40-60m	0	0	0	0	0	2	2	0	2	2	3	1	0	0	1	1	0	0	0	0
WCR12	60-80m	0	0	0	0	0	2	2	0	3	2	2	1	0	0	0	0	0	1	1	0
WCR12	80-100m	0	0	0	0	0	1	2	0	3	2	2	1	0	0	0	0	0	5	0	0
WCR12	100-120m	0	0	0	0	0	2	2	0	2	2	2	2	0	0	0	0	1	0	0	0
WCR12	120-140m	0	0	0	0	0	2	3	0	2	1	2	1	0	0	1	0	0	1	0	0
WCR12	140-160m	0	0	0	0	0	2	2	0	2	2	2	1	0	0	1	0	0	0	0	1
WCR12	160-180m	0	0	0	0	0	2	2	0	2	2	2	1	0	0	1	0	0	0	0	0
WCR12	180-200m	0	0	0	0	0	2	3	0	2	1	3	1	0	0	1	1	0	1	1	0
WCR12	200-220m	0	0	0	0	0	2	1	0	2	1	2	2	*	1	0	0	0	0	2	0
WCR12	220-240m	0	0	0	0	0	1	3	0	2	1	1	1	0	1	1	1	0	1	1	1
WCR12	240-250m	0	0	0	0	0	1	3	0	2	1	1	1	0	0	1	0	0	1	0	0
WCR12	average	0	0	0	0	0	2	2	0	2	2	2	1	*	*	8	3	2	11	5	2
WCR13	0-20m	0	0	0	0	0	3	2	0	2	2	2	2	0	1	0	0	0	0	0	0
WCR13	20-40m	0	0	0	0	*	2	1	0	2	2	2	2	0	0	0	0	0	1	0	0
WCR13	40-60m	0	0	0	0	0	2	1	0	3	1	2	1	0	*	0	0	0	0	1	0
WCR13	60-80m	0	0	0	0	0	1	1	0	3	1	2	1	0	0	0	0	0	2	0	0
WCR13	80-100m	0	0	0	0	0	2	1	0	2	1	3	2	0	0	1	0	1	0	1	1
WCR13	100-120m	0	0	*	0	0	2	1	0	2	1	2	1	0	0	0	0	0	2	1	0
WCR13	120-140m	0	0	0	0	0	1	1	0	2	1	2	1	0	0	0	1	0	0	0	0
WCR13	140-160m	0	0	0	0	0	2	1	0	2	1	3	1	0	0	0	0	0	2	0	0
WCR13	160-180m	0	0	0	0	0	2	1	0	2	1	3	1	0	0	1	1	0	1	1	0
WCR13	180-200m	0	0	0	0	0	2	1	0	2	1	3	1	0	0	3	0	0	4	2	0
WCR13	200-220m	0	0	0	0	*	2	1	0	3	1	2	1	0	*	1	0	0	0	1	0
WCR13	220-240m	0	0	0	0	0	2	1	0	3	1	2	1	0	0	2	0	0	0	0	0
WCR13	240-250m	0	0	0	0	0	1	1	0	3	1	2	1	0	0	0	0	0	4	1	0
WCR13	average	0	0	*	0	*	2	1	0	2	1	2	1	0	*	8	2	1	16	8	1
WCR14	0-20m	0	0	0	0	0	1	2	0	2	1	3	1	0	0	0	0	0	0	0	0
WCR14	20-40m	0	0	0	0	0	1	1	0	3	1	2	1	0	0	0	0	0	0	0	0
WCR14	40-60m	0	0	0	0	0	1	1	0	2	1	3	1	0	0	0	0	0	0	0	0

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR14	60-80m	*	0	0	0	*	1	2	0	2	1	4	1	0	0	1	1	0	0	0	0
WCR14	80-100m	*	0	0	0	0	1	1	0	2	1	4	1	0	0	1	0	0	0	0	0
WCR14	100-120m	0	0	0	0	0	1	1	0	3	1	3	1	0	0	0	0	0	1	0	0
WCR14	120-140m	0	0	0	0	*	1	1	*	2	1	4	1	0	0	1	1	0	0	0	0
WCR14	140-160m	0	0	0	0	*	2	1	0	2	1	4	1	0	0	0	0	0	0	2	0
WCR14	160-180m	0	0	0	0	*	3	1	0	2	2	4	1	0	0	0	0	0	0	0	0
WCR14	180-200m	0	0	0	0	0	2	2	0	2	1	3	1	0	0	1	0	0	0	0	0
WCR14	200-220m	0	0	0	0	1	1	1	0	1	2	4	1	0	0	1	0	0	0	0	0
WCR14	220-240m	0	0	0	0	1	2	1	*	1	2	4	1	0	0	1	0	0	2	1	0
WCR14	240-250m	0	0	0	0	1	1	1	*	1	2	3	1	0	0	3	0	0	0	0	0
WCR14	average	*	0	0	0	1	1	1	*	2	1	3	1	0	0	9	2	0	3	3	0
WCR15	0-20m	0	0	0	0	0	*	1	0	4	0	*	1	0	0	0	0	0	2	0	0
WCR15	20-40m	0	0	0	0	0	1	1	0	3	0	*	1	0	0	3	0	0	3	0	0
WCR15	40-60m	0	0	0	0	0	2	2	0	3	0	2	1	0	0	0	1	0	0	0	0
WCR15	60-80m	0	0	0	0	0	1	2	0	3	1	2	1	0	0	0	0	0	2	0	0
WCR15	80-100m	0	0	0	0	0	2	2	0	3	1	2	1	0	0	0	0	0	0	0	0
WCR15	100-120m	0	0	0	0	0	2	2	0	3	0	1	1	0	0	1	0	0	0	0	0
WCR15	120-140m	*	0	0	0	0	1	2	0	3	0	1	1	0	0	1	0	0	0	0	0
WCR15	140-160m	*	*	0	0	0	2	2	0	3	0	1	1	0	0	1	0	0	2	0	0
WCR15	160-180m	*	0	0	0	0	1	2	0	3	0	1	1	0	0	2	0	0	0	0	0
WCR15	180-200m	0	0	0	0	0	1	1	0	4	0	*	1	0	0	2	1	0	1	0	0
WCR15	200-220m	*	*	0	0	0	1	2	0	4	*	1	1	0	0	2	0	0	0	0	0
WCR15	220-240m	*	1	0	0	0	1	2	0	3	1	*	1	0	0	1	0	0	0	0	0
WCR15	240-250m	*	*	0	0	0	*	1	0	4	0	*	1	0	0	0	0	0	0	0	0
WCR15	average	1	1	0	0	0	1	2	0	3	1	1	1	0	0	13	2	0	10	0	0
WCR16	0-20m	0	0	0	0	0	1	1	*	3	3	2	1	0	*	3	1	0	4	1	0
WCR16	20-40m	0	0	0	0	0	1	1	0	3	1	1	1	0	*	0	0	1	2	0	0
WCR16	40-60m	0	0	0	0	0	1	1	*	3	2	1	1	0	1	0	0	0	3	1	0
WCR16	60-80m	0	0	0	0	0	1	1	*	3	1	1	1	0	0	0	0	0	3	0	0
WCR16	80-100m	0	0	0	0	0	1	1	0	3	1	1	1	0	0	2	0	0	2	2	0
WCR16	100-120m	0	0	0	0	0	1	1	0	3	1	1	1	0	0	0	2	0	0	1	0
WCR16	120-140m	0	0	0	0	0	1	1	0	3	1	1	1	0	0	2	0	0	2	0	0
WCR16	140-160m	*	0	0	0	0	1	1	0	3	2	1	2	0	0	0	2	0	2	0	0
WCR16	160-180m	0	0	0	0	0	1	1	0	3	1	1	1	0	0	0	0	0	1	0	0
WCR16	180-200m	0	0	0	0	0	1	1	0	3	1	1	1	0	0	1	0	0	1	0	0
WCR16	200-220m	*	0	0	0	0	1	1	0	3	1	1	1	0	0	2	0	0	0	1	0
WCR16	220-240m	*	0	0	*	0	1	1	0	3	1	1	1	0	0	0	2	0	0	0	0
WCR16	240-250m	*	0	0	0	0	1	1	0	3	*	1	1	*	0	1	0	0	0	0	0

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR16	average	*	0	0	*	0	1	1	*	3	1	1	1	*	*	11	7	1	20	6	0
WCR17	0-20m	0	0	0	0	0	1	1	0	4	2	1	*	0	0	3	1	0	0	1	0
WCR17	20-40m	0	0	0	0	0	1	1	0	4	2	1	1	0	0	4	0	1	0	0	0
WCR17	40-60m	0	0	0	0	0	1	1	0	4	1	1	1	0	0	1	1	0	1	0	0
WCR17	60-80m	0	0	0	0	0	1	1	0	4	1	1	1	0	0	0	0	1	0	0	2
WCR17	80-100m	0	0	0	0	0	1	1	0	4	2	1	*	0	0	0	2	0	2	1	0
WCR17	100-120m	0	0	0	0	0	1	*	0	3	2	*	2	0	*	0	0	1	1	0	1
WCR17	120-140m	0	0	0	0	0	1	*	0	3	1	*	2	0	2	3	1	0	1	0	1
WCR17	140-160m	0	0	0	0	0	1	*	0	4	2	*	1	0	0	2	0	0	0	0	0
WCR17	160-180m	0	0	0	0	0	1	1	0	4	2	*	*	0	0	4	0	0	0	0	0
WCR17	180-200m	0	0	0	*	0	*	*	0	4	2	*	*	0	0	1	1	0	0	0	0
WCR17	200-220m	0	0	0	0	0	1	1	0	4	1	*	*	0	0	1	0	0	0	1	0
WCR17	220-240m	0	0	0	0	0	1	1	0	4	1	1	*	0	0	1	0	0	1	1	0
WCR17	240-250m	0	0	0	0	0	1	1	0	4	1	1	*	0	0	1	1	0	0	1	0
WCR17	average	0	0	0	*	0	1	1	0	4	2	1	1	0	1	21	7	3	6	5	4
WCR18	0-20m	0	0	0	0	0	1	1	0	2	1	3	1	0	0	0	0	0	1	3	0
WCR18	20-40m	0	0	0	0	0	1	1	0	1	1	4	2	0	0	2	0	0	3	0	0
WCR18	40-60m	0	0	0	0	0	1	1	0	2	2	3	1	0	0	4	1	0	3	1	0
WCR18	60-80m	0	0	0	0	0	1	1	0	3	2	3	1	0	0	2	1	0	3	2	0
WCR18	80-100m	0	0	0	*	0	1	1	*	2	3	2	1	0	0	1	0	0	1	0	0
WCR18	100-120m	0	0	0	0	0	1	1	0	2	3	2	1	0	*	0	2	0	0	1	0
WCR18	120-140m	0	0	0	0	0	2	1	0	1	1	4	2	0	0	3	1	0	0	1	1
WCR18	140-160m	0	0	0	0	0	2	1	0	2	2	3	2	0	0	0	0	0	0	0	0
WCR18	160-180m	0	0	0	0	0	2	1	0	2	2	3	2	0	0	0	0	1	1	0	1
WCR18	180-200m	0	0	0	0	0	2	1	0	2	1	4	2	0	0	0	1	0	0	2	1
WCR18	200-220m	*	0	0	0	0	1	2	0	2	1	3	3	*	0	0	0	0	0	0	0
WCR18	220-240m	0	0	0	0	0	1	1	0	2	2	3	3	0	0	2	1	0	2	0	0
WCR18	240-250m	*	0	0	0	0	*	1	0	1	1	3	2	0	0	0	0	0	0	0	0
WCR18	average	*	0	0	*	0	1	1	*	2	2	3	2	*	*	14	7	1	14	10	3
WCR19	0-20m	*	*	0	0	0	*	*	0	1	1	3	3	0	0	0	0	0	0	0	0
WCR19	20-40m	0	*	0	0	0	2	0	0	1	2	3	3	0	0	2	0	1	2	0	0
WCR19	40-60m	0	1	0	0	0	2	0	0	1	2	3	3	0	0	0	0	0	0	0	0
WCR19	60-80m	0	*	0	0	0	1	*	0	2	1	2	2	0	0	0	0	0	0	2	0
WCR19	80-100m	*	1	0	0	0	*	*	0	2	1	2	2	0	0	0	0	0	0	1	1
WCR19	100-120m	0	0	0	0	0	*	*	0	1	2	3	0	0	0	1	0	0	0	0	0
WCR19	120-140m	0	0	0	0	0	1	*	0	2	1	3	2	0	0	1	0	0	0	0	1
WCR19	140-160m	0	0	0	0	0	*	*	0	2	1	2	1	0	0	6	0	0	0	4	0
WCR19	160-180m	0	0	0	0	0	*	0	0	3	1	2	1	0	0	3	1	0	2	0	0

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR19	180-200m	*	*	1	0	0	1	1	0	2	2	2	*	0	0	3	0	0	0	0	1
WCR19	200-220m	0	0	1	0	0	0	1	0	2	2	*	0	0	0	1	0	1	0	0	2
WCR19	220-240m	*	*	0	0	0	*	0	0	2	4	2	*	0	0	2	0	0	0	0	0
WCR19	240-250m	0	*	0	0	0	*	0	0	2	1	3	2	0	0	2	0	0	2	0	0
WCR19	average	*	1	*	0	0	1	1	0	2	2	2	2	0	0	21	1	2	6	7	5
WCR20	0-20m	0	*	*	0	0	*	0	0	3	1	1	1	0	0	1	0	0	0	0	0
WCR20	20-40m	0	*	0	0	0	*	0	0	2	1	2	2	0	0	1	0	0	0	1	0
WCR20	40-60m	0	1	*	0	0	1	0	0	3	2	2	2	0	0	0	0	2	1	1	0
WCR20	60-80m	0	1	0	0	0	1	0	0	3	1	1	1	0	0	0	0	0	2	0	0
WCR20	80-100m	0	2	0	0	0	2	0	0	3	1	1	1	0	0	1	0	0	1	0	0
WCR20	100-120m	0	2	0	0	0	2	0	0	3	1	1	1	0	0	0	0	0	0	0	0
WCR20	120-140m	0	2	0	0	0	2	0	0	3	1	1	1	0	0	1	0	0	0	0	0
WCR20	140-160m	0	2	0	0	0	2	1	1	3	1	1	2	0	0	0	0	1	0	2	0
WCR20	160-180m	0	*	0	0	0	*	1	1	2	2	2	2	0	0	1	0	1	0	1	0
WCR20	180-200m	0	0	*	0	0	1	1	1	2	3	1	1	0	0	1	0	0	0	0	0
WCR20	200-220m	0	*	*	0	0	*	1	1	2	3	1	1	0	0	1	1	0	0	0	0
WCR20	220-240m	0	0	*	0	0	1	1	1	2	3	1	1	0	0	2	0	0	0	0	0
WCR20	240-250m	0	*	*	0	0	1	1	1	2	3	1	1	0	0	0	0	0	1	0	0
WCR20	average	0	1	1	0	0	1	1	1	2	2	1	1	0	0	9	1	4	5	5	0
WCR21	0-20m	0	1	0	0	0	0	0	0	4	0	*	*	0	0	1	0	1	0	0	0
WCR21	20-40m	0	2	0	0	0	0	0	0	3	0	2	1	0	0	0	0	0	1	1	0
WCR21	40-60m	0	2	0	0	0	0	0	0	3	0	2	*	0	0	1	0	1	0	1	0
WCR21	60-80m	0	2	0	0	0	0	0	0	4	0	2	1	0	0	0	0	0	3	0	0
WCR21	80-100m	0	2	0	0	0	0	0	0	4	0	*	*	0	0	0	0	0	2	0	0
WCR21	100-120m	0	2	0	0	0	0	0	0	4	*	*	1	0	0	0	0	0	0	0	0
WCR21	120-140m	0	2	0	0	0	0	0	0	4	*	*	1	0	0	0	0	1	2	0	0
WCR21	140-160m	0	1	0	0	0	0	0	0	3	1	1	*	0	0	0	3	0	2	0	0
WCR21	160-180m	0	1	*	0	0	1	*	0	2	1	2	1	0	0	2	0	0	0	0	0
WCR21	180-200m	0	0	*	0	0	1	*	0	2	2	2	0	0	0	0	0	0	1	0	0
WCR21	200-220m	0	0	*	0	0	2	1	0	1	2	3	2	0	0	0	0	1	0	0	0
WCR21	220-240m	0	0	0	0	0	0	0	0	2	4	3	*	0	1	0	0	1	0	0	0
WCR21	240-250m	0	*	0	0	0	*	0	0	2	3	3	0	0	1	0	0	0	1	0	0
WCR21	average	0	1	*	0	0	1	*	0	3	2	2	1	0	*	4	3	5	12	2	0
WCR22	0-20m	0	*	0	0	0	0	0	0	3	1	1	2	0	0	0	0	0	0	0	0
WCR22	20-40m	0	1	0	0	0	0	0	0	4	1	1	1	0	0	0	0	0	1	0	0
WCR22	40-60m	0	1	0	0	0	0	0	0	5	2	1	1	0	0	1	1	0	1	0	0
WCR22	60-80m	0	1	0	0	0	0	0	0	5	1	1	1	0	0	1	0	0	0	0	0
WCR22	80-100m	0	*	0	0	0	0	0	0	5	1	1	1	0	0	0	1	0	0	0	0

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR22	100-120m	0	1	0	0	0	0	*	0	5	0	1	1	0	0	1	0	0	2	0	0
WCR22	120-140m	0	1	0	*	0	0	0	0	5	0	1	1	0	0	0	0	0	2	0	0
WCR22	140-160m	0	*	0	0	0	*	0	0	5	0	1	1	0	0	0	1	0	1	0	0
WCR22	160-180m	0	1	0	0	0	*	0	0	5	*	1	1	0	0	0	0	0	0	0	0
WCR22	180-200m	0	1	0	0	0	0	0	0	5	1	1	1	0	0	1	0	0	0	0	0
WCR22	200-220m	0	1	0	0	0	0	0	0	5	0	1	1	0	0	0	0	0	1	0	0
WCR22	220-240m	0	*	0	0	0	0	0	0	5	1	1	1	0	0	0	0	0	0	0	0
WCR22	240-250m	0	0	0	0	0	*	0	0	4	1	1	2	0	0	0	0	0	0	0	0
WCR22	average	0	1	0	*	0	*	*	0	5	1	1	1	0	0	4	3	0	8	0	0
WCR23	0-20m	0	0	1	0	0	0	0	0	1	1	4	3	0	*	1	0	0	1	0	0
WCR23	20-40m	0	0	1	0	0	1	1	0	2	1	2	1	0	1	0	0	0	0	0	0
WCR23	40-60m	0	1	0	0	0	1	2	0	3	1	2	*	0	0	0	0	0	0	0	0
WCR23	60-80m	0	*	*	0	0	1	2	0	3	1	2	1	0	0	0	0	0	1	0	0
WCR23	80-100m	0	1	0	0	0	1	2	0	3	1	3	1	0	0	0	0	1	0	0	0
WCR23	100-120m	0	1	*	0	0	1	2	0	2	1	4	1	0	0	0	0	0	0	0	0
WCR23	120-140m	0	1	0	0	0	1	1	0	2	1	3	1	0	0	1	0	0	2	0	0
WCR23	140-160m	0	1	0	0	0	*	1	0	3	1	2	1	0	0	2	0	0	0	0	0
WCR23	160-180m	0	2	0	0	0	0	1	0	3	2	2	1	0	0	0	0	0	0	0	0
WCR23	180-200m	0	1	0	0	0	*	1	0	3	2	2	1	0	0	0	0	0	0	0	0
WCR23	200-220m	0	*	*	0	0	*	1	0	4	2	2	1	0	0	2	0	0	0	1	0
WCR23	220-240m	0	*	0	0	0	*	2	0	4	1	2	1	0	0	0	0	0	0	0	0
WCR23	240-250m	0	1	0	0	0	*	1	0	4	1	2	1	0	0	0	0	0	0	0	0
WCR23	average	0	1	1	0	0	1	1	0	3	1	2	1	0	*	6	0	1	4	1	0
WCR24	0-20m	0	0	2	0	0	*	1	*	2	2	2	1	0	0	0	0	0	2	1	0
WCR24	20-40m	0	0	*	0	0	0	*	*	2	2	2	*	0	0	0	0	0	1	1	0
WCR24	40-60m	0	0	*	0	0	*	1	0	3	2	2	*	0	0	1	0	0	1	0	0
WCR24	60-80m	0	0	0	0	0	*	1	*	3	3	2	1	0	0	1	0	0	0	1	0
WCR24	80-100m	0	0	0	0	0	1	*	*	3	2	2	2	0	0	1	0	0	1	0	0
WCR24	100-120m	0	0	*	0	0	1	*	*	1	2	2	4	0	*	1	0	0	0	0	0
WCR24	120-140m	0	0	0	0	0	*	*	0	2	1	2	4	0	0	0	0	0	0	1	0
WCR24	140-160m	0	0	*	0	0	*	0	0	2	1	2	4	0	0	0	0	0	0	0	0
WCR24	160-180m	0	0	2	0	0	*	*	*	2	*	2	3	0	0	0	0	0	0	0	0
WCR24	180-200m	0	0	*	0	0	*	*	0	2	1	2	2	0	0	1	0	0	0	0	0
WCR24	200-220m	0	0	0	0	0	1	*	0	2	1	2	2	0	0	2	0	0	0	0	0
WCR24	220-240m	0	0	0	0	0	1	1	*	2	2	3	2	0	1	1	0	0	0	0	0
WCR24	240-250m	0	0	1	0	0	*	0	0	1	1	1	0	0	2	0	0	0	0	0	0
WCR24	average	0	0	1	0	0	1	1	1	2	2	2	2	0	1	8	0	0	5	4	0
WCR25	0-20m	0	0	0	0	*	1	1	0	2	2	3	1	0	0	0	0	0	0	0	0

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR25	20-40m	0	0	0	0	*	1	1	0	2	2	3	1	0	0	0	0	0	2	0	0
WCR25	40-60m	0	0	0	0	0	*	0	0	2	2	3	2	0	0	0	0	0	1	1	0
WCR25	60-80m	0	0	0	0	*	1	0	0	3	2	2	1	0	1	0	2	0	0	1	0
WCR25	80-100m	0	0	0	0	*	1	1	0	3	2	2	2	0	1	2	0	0	0	2	0
WCR25	100-120m	0	0	0	0	2	2	1	1	2	2	2	2	*	1	2	0	0	4	0	0
WCR25	120-140m	0	0	0	0	2	2	1	1	2	1	2	2	*	2	1	1	1	0	1	0
WCR25	140-160m	0	0	0	0	1	2	1	1	2	1	2	2	*	1	0	1	0	0	0	0
WCR25	160-180m	0	0	0	0	*	1	1	1	3	1	2	1	0	0	1	1	1	0	0	0
WCR25	180-200m	0	0	0	0	*	2	1	0	2	1	2	2	0	1	0	0	0	0	2	2
WCR25	200-220m	0	0	0	0	*	2	1	*	1	1	3	2	0	1	1	0	0	0	0	0
WCR25	220-240m	0	0	0	0	0	1	1	0	2	1	4	1	0	0	0	0	0	1	2	0
WCR25	240-250m	0	0	0	0	*	1	1	0	3	1	3	1	0	0	0	0	0	0	0	0
WCR25	average	0	0	0	0	1	1	1	1	2	1	2	2	*	1	7	5	2	8	9	2
WCR26	0-20m	0	1	0	0	0	*	1	0	3	3	2	1	0	0	0	0	0	1	1	0
WCR26	20-40m	0	1	0	0	0	0	1	0	3	3	2	1	0	0	0	0	0	1	1	0
WCR26	40-60m	0	1	0	0	0	0	1	0	3	2	2	1	0	0	1	0	0	1	1	0
WCR26	60-80m	0	1	*	0	0	0	1	0	4	1	1	1	0	0	1	0	0	0	0	0
WCR26	80-100m	0	1	0	0	0	1	1	*	3	2	1	1	0	0	0	0	0	2	2	0
WCR26	100-120m	0	1	0	0	0	1	1	1	3	1	1	2	0	0	2	0	0	1	0	0
WCR26	120-140m	0	1	0	0	0	1	1	1	3	2	1	1	0	0	1	0	0	0	0	0
WCR26	140-160m	0	1	0	0	0	0	*	*	3	2	1	1	0	0	1	0	0	0	0	0
WCR26	160-180m	0	2	0	0	0	*	*	*	3	2	2	1	0	0	1	2	0	0	0	1
WCR26	180-200m	0	1	0	0	0	0	1	1	4	1	1	1	0	0	2	0	0	1	1	0
WCR26	200-220m	0	2	0	0	0	0	*	*	3	2	1	1	0	0	0	0	0	1	0	1
WCR26	220-240m	0	0	0	0	0	2	1	2	3	2	1	2	0	0	0	1	0	1	0	0
WCR26	240-250m	0	*	0	0	0	2	2	2	2	1	1	2	0	*	0	1	0	1	0	0
WCR26	average	0	1	*	0	0	1	1	1	3	2	1	1	0	*	9	4	0	10	6	2
WCR27	0-20m	0	0	0	0	0	2	3	0	3	2	1	*	0	0	0	0	0	2	0	0
WCR27	20-40m	0	0	0	0	0	2	2	0	3	2	1	1	0	0	0	0	0	1	0	0
WCR27	40-60m	0	0	0	0	0	1	2	0	3	2	2	1	0	0	0	0	0	0	0	0
WCR27	60-80m	0	0	0	0	0	3	2	0	2	2	3	1	0	0	0	0	0	0	0	0
WCR27	80-100m	0	0	0	0	0	1	2	0	3	2	2	1	0	0	0	0	0	1	0	0
WCR27	100-120m	0	0	0	0	0	1	2	0	4	1	2	1	0	0	0	0	0	2	0	0
WCR27	120-140m	0	0	0	0	0	2	2	0	3	*	2	2	0	0	0	0	0	0	0	0
WCR27	140-160m	0	0	0	0	0	1	1	0	3	0	2	3	0	*	0	0	0	0	0	0
WCR27	160-180m	0	0	0	0	0	2	1	0	1	0	2	3	0	0	0	1	0	0	0	0
WCR27	180-200m	0	0	0	0	0	*	1	0	1	0	2	4	0	1	2	0	0	2	0	0
WCR27	200-220m	0	0	*	0	0	*	1	0	1	1	1	4	0	*	1	0	0	1	0	0

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR27	220-240m	0	*	*	0	0	1	2	0	1	1	2	4	0	1	2	0	0	1	0	0
WCR27	240-250m	0	*	0	0	0	*	*	*	1	2	3	3	0	*	0	0	0	0	0	0
WCR27	average	0	*	*	0	0	1	2	*	2	1	2	3	0	1	5	1	0	10	0	0
WCR28	0-20m	0	0	0	*	0	0	*	0	4	1	1	1	0	0	3	3	0	2	2	0
WCR28	20-40m	0	0	0	*	0	1	*	0	4	1	1	1	0	0	0	0	0	0	0	0
WCR28	40-60m	0	0	0	0	*	2	*	0	4	1	1	1	0	0	0	0	0	0	0	0
WCR28	60-80m	0	0	0	0	0	2	*	0	4	2	2	*	0	0	2	0	0	1	0	0
WCR28	80-100m	0	0	0	0	*	2	*	*	3	2	2	1	1	0	0	0	0	1	0	0
WCR28	100-120m	0	0	0	0	1	1	*	*	3	2	1	0	0	0	1	0	0	0	0	0
WCR28	120-140m	0	0	0	0	1	1	*	*	3	1	1	1	0	0	0	0	0	1	1	0
WCR28	140-160m	0	0	0	0	*	2	*	0	3	2	2	1	1	*	0	0	0	0	0	0
WCR28	160-180m	0	0	0	0	2	*	*	0	3	2	1	*	2	1	2	0	0	2	0	0
WCR28	180-200m	0	0	0	0	*	*	*	0	3	2	*	1	0	0	5	0	0	2	1	0
WCR28	200-220m	0	0	*	0	2	1	*	0	3	2	*	*	0	0	0	0	0	2	0	0
WCR28	220-240m	0	0	*	0	1	2	*	0	3	2	*	*	0	*	0	1	0	3	1	0
WCR28	240-250m	0	0	1	0	3	*	1	1	3	2	1	*	0	0	1	0	0	0	0	1
WCR28	average	0	0	*	*	1	1	1	1	3	2	1	1	1	*	14	4	0	14	5	1
WCR29	0-20m	0	0	*	0	*	3	2	2	1	2	2	1	0	1	0	0	0	0	0	0
WCR29	20-40m	0	0	*	0	0	3	2	1	1	2	3	2	0	*	0	0	0	1	0	0
WCR29	40-60m	0	0	0	0	0	2	2	1	2	2	2	2	0	*	0	0	0	0	0	0
WCR29	60-80m	0	0	0	0	0	2	1	0	2	2	2	2	0	0	1	0	0	0	0	0
WCR29	80-100m	0	0	0	0	0	2	1	0	3	2	3	2	0	0	0	0	0	0	0	0
WCR29	100-120m	0	0	0	0	0	1	1	0	4	2	2	2	0	0	1	0	0	0	1	0
WCR29	120-140m	0	0	0	0	0	2	1	0	3	2	2	2	0	0	1	0	0	0	0	0
WCR29	140-160m	0	0	0	0	0	2	2	1	3	2	2	3	0	0	1	0	0	1	0	0
WCR29	160-180m	0	0	0	0	0	3	2	0	2	1	2	4	0	0	0	0	0	0	0	0
WCR29	180-200m	0	0	0	0	0	3	2	0	2	1	2	3	0	0	0	0	0	0	0	0
WCR29	200-220m	0	0	*	0	0	2	2	0	2	*	2	3	0	0	1	0	0	0	0	0
WCR29	220-240m	0	0	0	0	0	2	2	0	2	1	2	3	0	0	0	0	0	1	0	0
WCR29	240-250m	0	0	0	0	0	2	1	0	2	*	2	3	0	0	0	0	0	0	0	0
WCR29	average	0	0	*	0	*	2	2	1	2	2	2	3	0	*	5	0	0	3	1	0
WCR30	0-20m	0	0	0	0	0	2	1	0	3	2	2	1	0	0	0	0	0	1	0	0
WCR30	20-40m	0	0	0	0	0	1	1	0	3	1	2	1	0	0	0	0	0	2	0	0
WCR30	40-60m	0	0	0	0	0	1	1	0	3	1	2	1	0	0	1	0	0	0	0	2
WCR30	60-80m	0	0	*	0	0	2	1	0	3	2	2	2	0	1	0	0	0	4	0	1
WCR30	80-100m	0	0	2	0	0	2	1	*	2	2	2	2	0	2	0	0	1	4	0	1
WCR30	100-120m	1	0	*	0	0	2	1	1	2	2	2	2	0	1	0	0	0	0	0	0
WCR30	120-140m	0	0	*	0	0	1	1	*	3	2	2	3	0	1	0	0	0	1	2	0

Site	Transect segment	Mistletoe	Silver Wattle	Blackwood	Cherry Ballart	Musk Daisy-bush	Shrubs > 1m	Shrubs < 1m	Coprosma	Leaf litter	Bracken	Native grasses	Rushes/sedges	Tree ferns	Ground ferns	Logs 40-59cm	Logs 60-79cm	Logs > 80cm	Stumps 40-59cm	Stumps 60-79cm	Stumps > 80cm
WCR30	140-160m	0	0	*	0	1	1	1	*	3	2	2	3	0	1	1	0	0	0	0	0
WCR30	160-180m	0	0	*	0	1	1	1	*	3	2	2	3	0	1	0	0	0	0	0	1
WCR30	180-200m	0	0	*	0	2	2	1	*	3	1	1	2	0	2	0	0	0	0	0	0
WCR30	200-220m	0	0	*	0	4	1	*	*	3	*	1	1	0	2	0	1	0	0	2	0
WCR30	220-240m	0	0	0	0	4	*	*	*	2	*	1	2	0	3	1	0	0	3	0	0
WCR30	240-250m	0	0	2	0	3	*	*	*	1	1	1	2	*	3	0	0	0	4	0	0
WCR30	average	*	0	1	0	1	1	1	1	3	2	2	2	*	2	3	1	1	19	4	5

